



Imgen BioSciences, Inc.

151 Martine Street, Fall River, MA 02723, USA Tel: 1 (781) 850-2047 www.imgenbiosciences.com

B-1012: Hygromycin B-Agarose 4B

Product Name:	Hygromycin B-Agarose 4B
Catalogue No:	B-1012
Antigen/Ligand:	Hygromycin B
Antigen/Ligand Concentration:	Not determined
Bead Structure:	4% agarose
Bead Size Range:	45-165 μm
Mean Bead Size:	90 μm
Linker:	6-aminohexanoic acid
Linker Space:	8 atoms
Size:	1 g
Form:	Lyophilized powder (stabilized with lactose and dextran)
Swelling:	1 g swells to 3-4 ml
Binding Capacity:	Antibodies: 15-20 mg/ml of drained gel
Max Linear Flow Rate*:	75 cm/h at 25°C, HR 16/10 column, 5 cm bed height
Storage Temp:	Keep at 2-8°C.
Applications:	Used as capture antigen/ligand for the separation or purification of antibodies specific to hygromycin B by affinity chromatography and/or spin down.

Brief description:

Hygromycin B is covalently conjugated to 4% beaded agarose. One to two amine groups in the hygromycin B is directly linked to the active group of N-hydroxysuccinimide on the pre-activated CH-agarose. This product is produced specifically for the binding of hygromycin B binding agents such as Hygromycin B-specific antibodies.

There is approximately 15 μmole of active hydroxycuccinidie group/ml of drained agarose 4B gel. For coupling the ligand/antigen, 1 volume of coupling buffer containing 100 mM Hygromycin B sodium is added to a same volume of swollen agarose 4B gel. Assuming 30% of active hydroxycuccinidie group is crosslinked with Hygromycin B that has 2 amine group per molecule, the final concentration of Hygromycin B is 4.5 $\mu\text{mole/ml}$ of drained agarose gel.

PLEASE note that this product is intended for research use only; not for diagnostic or clinical use.

*Linear flow rate (cm/hr) = volumetric flow rate (cm^3/min) X 60min/Cross sectional area of column (cm^2)

(Updated April, 2013)